

Entanglement, quantum phase transitions, and density matrix renormalization

Tobias J. Osborne^{1,2,*} and Michael A. Nielsen^{2,†}

¹*Department of Mathematics, University of Queensland 4072, Australia*

²*Centre for Quantum Computer Technology, University of Queensland 4072, Australia*

(Dated: February 1, 2008)

We investigate the role of entanglement in quantum phase transitions, and show that the success of the density matrix renormalization group (DMRG) in understanding such phase transitions is due to the way it preserves entanglement under renormalization. We provide a *reinterpretation* of the DMRG in terms of the language and tools of quantum information science which allows us to rederive the DMRG in a physically transparent way. Motivated by our reinterpretation we suggest a modification of the DMRG which manifestly takes account of the entanglement in a quantum system. This modified renormalization scheme is shown, in certain special cases, to preserve more entanglement in a quantum system than traditional numerical renormalization methods.

PACS numbers: 03.65.Ud, 73.43.Nq, 05.10.-a

A fundamental difference between quantum and classical physics is the possible existence of nonclassical correlations between distinct quantum systems. The physical property responsible for the nonclassical correlations is called *entanglement* [1, 2, 3]. Recently [4, 5, 6], it has been conjectured that entanglement plays a critical role in the quantum phase transitions which occur in interacting quantum lattice systems [7] at zero temperature.

Unlike ordinary phase transitions, quantum phase transitions take place at zero temperature as one or more physical parameters of the system are varied. At the critical point there are long-range correlations in the system, as in a conventional phase transition. However, because the system is at zero temperature, and assuming there is no ground-state degeneracy, the system must be in a pure state. It follows that the correlations which are the principal experimental signature of a quantum phase transition are due to long-range entanglement in the ground state of the system.

A variety of methods have been developed to understand quantum phase transitions. One of the most successful is the density matrix renormalization group (DMRG) [8], a numerical technique for finding accurate approximations to the ground and low-lying excited states of interacting quantum lattice systems. The DMRG was developed in response to the poor results obtained from Wilson's numerical renormalization group (RG) [9] when applied to general quantum lattice systems. White [10] identified the mathematical origin of the breakdown of Wilson's RG as the failure of the method to properly take into account the effect of boundary conditions during iterations.

The focus of this paper is not on proving technical results. Rather, the purpose is to provide a novel physical picture of the DMRG and quantum phase transitions based on the theory of entanglement developed by quantum information science [11, 12]. This physical picture provides a fruitful way of interpreting the properties of condensed-matter systems, substantially simplifies and

physically motivates existing derivations of properties of the DMRG, and motivates us to propose renormalization schemes extending the DMRG. We will explain the physical origin of the breakdown of Wilson's RG as a failure to take into account the effects of quantum entanglement in the ground state of the system, and show that the DMRG can be viewed as a natural extension of Wilson's RG to take into account the contribution entanglement makes to correlations in the ground state. Unfortunately, owing to the limited results available on entanglement measures, our proposed modifications to the DMRG cannot be derived in closed form. For this reason we are forced to implement the modified renormalization scheme numerically for idealised example systems with only small numbers of spins.

The structure of the paper is as follows. First, we explain the critical role entanglement plays in any long-range correlations present in a quantum system at zero temperature. Second, the physical origin of the failure of Wilson's RG is identified. The relationship between the DMRG and its treatment of entanglement is then elucidated. Finally, a numerical renormalization scheme is presented which has the property of preserving more entanglement in a quantum system than the DMRG. The differences in the entanglement preserved by the various renormalization schemes is illustrated using the 1D Heisenberg model.

We now consider the physical origin of the correlations which occur in systems exhibiting a quantum phase transition. For concreteness, we restrict attention to a lattice of spin- $\frac{1}{2}$ particles. If the ground state were a product state, then a simple calculation shows that the spin-spin correlation function $\langle \sigma_i^\alpha \sigma_j^\beta \rangle - \langle \sigma_i^\alpha \rangle \langle \sigma_j^\beta \rangle$ is identically zero. Thus, if the correlation function is non-zero then the ground state must be entangled, and large values of the correlation function imply a highly entangled ground state. It is well known that for general quantum lattice systems the correlation function decays exponentially as a function of the separation $|i - j|$ when the system is far

from criticality [7]. When the system is at a critical point, the correlations decay only as a polynomial function of the separation. At this point a fundamental change in the ground state has occurred. The physical property responsible for these quantum correlations is entanglement, so that for systems approaching a critical point the structure of the entanglement in the ground state must undergo a transition. Evidence for such a transition has been obtained for the transverse Ising model [18].

The standard technique used to derive the polynomial decay of correlations at the critical point is the *renormalization group* [9]. Renormalization group techniques as applied to quantum field theory and quantum lattice systems are now well developed (see, for example, [13, 14], for reviews). The renormalization group works by successively discarding degrees of freedom in a system until the dominant terms may be identified. Each renormalization scheme makes a decision about what degrees of freedom in a system are important and this decision, in turn, dictates which class of systems the method is best suited to. Prior to 1992 the success of the numerical implementation of the renormalization group for quantum systems was limited to the Kondo problem [9], and attempts to apply Wilson's RG to other quantum lattice systems suffered from poor convergence and low accuracy.

Briefly, Wilson's numerical RG procedure works by isolating a portion A of a quantum lattice system containing L sites and exactly diagonalising the Hamiltonian H_L of the system restricted to these sites. The subspace W of the m lowest eigenvalues and eigenvectors is obtained. (Note that m is a parameter we choose. It quantifies how many degrees of freedom we keep when renormalizing.) For example, for system A we might pick two adjacent spins on the lattice, and then determine the two-dimensional subspace W corresponding to the two lowest eigenvalues of H_L . Once this is done all operators are "renormalized" (effectively, averaging out degrees of freedom thought to be unimportant), by projecting onto the m -dimensional subspace W . This allows us to renormalize the entire lattice, replacing the two spins originally in A with a single renormalized spin. On the new renormalized lattice a site is added to A and the interactions with the new site are taken into account by constructing a new Hamiltonian H_{L+1} . The exact diagonalisation is performed again, keeping the m lowest eigenvalues and eigenvectors and the subspace W is updated. The expectation value of any operator of interest is calculated with respect to the ground state of A (which is the approximation to the overall ground state), and as the procedure is repeated the change in these expectation values are noted. Convergence is deemed to have been obtained when this difference becomes smaller than some prespecified value.

Wilson's RG procedure was expected to work because it was assumed that the only degrees of freedom contributing significantly to the overall ground state would

be the ground state and low-energy excitations of the lattice portion A . This assumption turns out not to be the case, and Wilson's RG works poorly for most systems. White [10] gave a mathematical explanation of why this is the case, in terms of the failure of Wilson's RG to properly take into account boundary conditions on the system being renormalized. Using the tools of quantum information science, we can provide another way to interpret the failure of Wilson's RG. The reason the assumption fails is that the ground state of an interacting lattice is typically entangled, and thus a subsystem of the total lattice cannot be assigned a definite state. Indeed, Wilson's RG ignores the terms coupling A to the remainder of the lattice, which are the very terms responsible for the entanglement in the ground state causing the correlations which signify a quantum phase transition.

The DMRG improves upon Wilson's scheme by taking into account the entanglement between system A and the remainder of the lattice. It does this by introducing a second subsystem of the lattice, B (the *environment*), in addition to the portion A (now referred to as the *principal system*) isolated by Wilson's RG procedure. For example, if the principal system A is two adjacent spins in a one-dimensional lattice, then B might consist of two spins, one on either side of A . A and B together are referred to as the *superblock*. Using the physical picture based upon entanglement, the DMRG may be described as follows. First, construct the superblock Hamiltonian and find its ground state. Of course, this ground state does not properly describe the entanglement between the superblock and the remainder of the lattice. However, provided B is chosen appropriately, most of the entanglement between A and the remainder of the lattice will in fact be entanglement between A and B . Physically, we expect this to happen if B contains all those sites to which A most strongly couples. The second step is to renormalize A by projecting onto those degrees of freedom of A which contribute most to the entangled ground state of the superblock, in a sense made more precise below. That is, we choose an m -dimensional subspace W of system A , and renormalize the operators for A in a similar manner to that described for Wilson's RG. The difference between the two approaches is in how W is chosen: in Wilson's RG, W is chosen without taking account of ground state entanglement, while in the DMRG ground state entanglement is critical. This method is iterated until convergence (using the same criteria as for Wilson's RG) is obtained.

Motivated by this physical picture, we can now explain the mathematics of the DMRG. Our description is equivalent to White's [10], but is presented using techniques of quantum information science which simplify the proof, and highlight the key physical role entanglement plays in the DMRG. The central concept we need is the *Schmidt decomposition* [11], one of the main tools used in the study of entangled states. The Schmidt decomposition

states that a pure state $|\Psi\rangle_{AB}$ of a *bipartite* system with components A and B may be written as a single sum over positive real co-efficients $\sqrt{p_\alpha}$,

$$|\Psi\rangle_{AB} = \sum_{\alpha} \sqrt{p_\alpha} |u^\alpha\rangle |v^\alpha\rangle, \quad (1)$$

where $|u^\alpha\rangle$ and $|v^\alpha\rangle$ are orthonormal eigenvectors of the reduced density matrices ρ_A and ρ_B , respectively. It is easy to show that $|\Psi\rangle$ is entangled if and only if there is more than one term in Eq. (1).

Which degrees of freedom should we keep to properly take into account the entanglement between A and B ? One reasonable choice is provided by the DMRG projection P , which is defined to be the m -dimensional projection on system A that minimizes the functional

$$\mathcal{S}(P) = ||\Psi\rangle_{AB} - P|\Psi\rangle_{AB}|^2. \quad (2)$$

The motivation for introducing this functional is to isolate the degrees of freedom which, if discarded, would change the state the least. To determine the projection which minimizes this functional we write the state $|\Psi\rangle$ of the superblock in Schmidt form, $|\Psi\rangle = \sum_{\alpha} \sqrt{p_\alpha} |u^\alpha\rangle |v^\alpha\rangle$. Simple algebra shows that

$$||\Psi\rangle_{AB} - P|\Psi\rangle_{AB}|^2 = 1 - \text{tr} \left(P \sum_{\alpha} p_\alpha |u^\alpha\rangle \langle u^\alpha| \right). \quad (3)$$

Thus minimizing the functional in Eq. (2) is equivalent to maximizing $\text{tr} (P \sum_{\alpha} p_\alpha |u^\alpha\rangle \langle u^\alpha|)$. The Ky Fan maximum principle [15] tells us that the maximum in this expression is achieved by choosing P to be the projector onto the space spanned by the $|u^\alpha\rangle$ with the m largest values p_α .

Summarizing, the physical origin for the failure of Wilson's RG and the success of the DMRG lies in the way each method deals with entanglement. Typically the ground state of a lattice system exhibits quantum correlations. In order to represent the state of an isolated portion of lattice sites A it is necessary to take into account the correlations between A and the rest of the lattice C . Due to the nonlocal nature of the entanglement responsible for the correlations, it is impossible to represent the state of AC using only the local degrees of freedom of A . Thus Wilson's RG cannot represent the correlations of A with C unless a full exact diagonalisation is performed. The DMRG approximates the correlations of A with C by introducing a subsystem B of C in such a way that the entanglement between A and B approximates the entanglement between A and C .

We now outline another technique to determine the best degrees of freedom to retain during the renormalization step that is easily generalized to mixed states. The best degrees of freedom to retain may be thought of as identifying which m -dimensional subspace will best preserve the overlap, or *fidelity*, of the original state with a

state that has support only on the smaller subspace. We write the renormalized state in the m -dimensional subspace in Schmidt form $|\tilde{\Psi}\rangle = \sum_{\beta=1}^m \sqrt{q_\beta} |w^\beta\rangle |x^\beta\rangle$. The quantity we wish to maximize is the fidelity between the renormalized and unrenormalized states, $\langle \tilde{\Psi} | \Psi \rangle$. This problem has been considered previously [16, 17] where it was shown that the maximum fidelity occurs when the Schmidt basis for $|\tilde{\Psi}\rangle$ corresponds to the Schmidt basis elements of $|\Psi\rangle$ with the m largest Schmidt coefficients. This derivation is therefore equivalent to the projection method because the same subspaces are preserved. The subspace-fidelity derivation has the added benefit of being physically transparent as well as being easily generalized to mixed states. Indeed, when the state of the superblock is mixed, that is, at finite temperature, the projection method may be used to show [8] that the optimal degrees of freedom to retain in the renormalization step are the m eigenvectors of the reduced density matrix for A with largest eigenvalues. This result may be verified with a subspace-fidelity argument where the expression for the fidelity of a pair of mixed density matrices is $F(\rho_1, \rho_2) = \text{tr} (\sqrt{\sqrt{\rho_1} \rho_2 \sqrt{\rho_1}})^2$.

The DMRG is an excellent numerical scheme for many applications because the state after the projection is usually close to the original. However, there is strong evidence that the DMRG does not reproduce the algebraic decay of correlations with respect to separation at the critical point [19], which is the standard signature of a quantum phase transition. We conjecture that the origin of this failure is due to the fact that while the DMRG does preserve *some* of the entanglement between A and the remainder of the lattice, it does not preserve the *maximal* amount of entanglement. To make this intuition more precise, we need a good quantitative measure of how much entanglement is present in a quantum state. For pure states such a measure of entanglement between A and B is the von Neumann entropy of the individual systems A and B alone, $S(A) = -\text{tr} \rho_A \log \rho_A = S(B)$ [3]. We conjecture that the reason the DMRG does not reproduce the algebraic decay of correlations at the critical point is because it minimizes the difference (2) rather than the difference between the entanglement of AB and the entanglement of AB after the projection.

This conjecture motivates the construction of a projection preserving more entanglement between the system and environment than the DMRG. For the case where the superblock is in a pure state we wish to determine which projection P on A when applied to the joint state $|\Psi\rangle_{AB}$ of AB will preserve the maximal amount of entanglement between A and B . This is equivalent to maximizing

$$\mathcal{F} = S(P\rho_A P / \text{tr}(P\rho_A)), \quad (4)$$

over all m -dimensional projections P on A . To do this maximization numerically, we introduce a basis $\{|\phi_i\rangle\}$ for A and write $P = \sum_i |\phi_i\rangle \langle \phi_i|$ where the sum runs

over some subset of the basis. The projection P is determined by varying the directions of the basis vectors until the quantity in (4) is maximized. Numerically this is done by introducing a Lagrange multiplier for each basis vector to enforce orthonormality. We refer to the projection P constructed by the procedure outlined here as the entanglement maximization projection (EMP).

For nonzero temperatures the state of the superblock ρ_{AB} is mixed. The derivation of the EMP outlined in the previous paragraph requires a measure of the entanglement of A with B for mixed states in order to maximize (4). For our calculations we use the best-understood measure of mixed-state entanglement, the *entanglement of formation* $E(A : B)$ [3] of the joint state ρ_{AB} . The construction of the EMP is otherwise unchanged. Unfortunately, no simple formula for the entanglement of formation has yet been obtained. For this reason we are only able to calculate the EMP for systems with small numbers of spins. When a simple formula for a good mixed-state entanglement measure is obtained it may be possible to derive a general formula for the EMP.

The EMP has the property that entanglement in the system is preserved maximally. The EMP does not minimize the functional Eq. (2) so that, in general, the state is quite different after the projection. For this reason the EMP is not appropriate for most calculations. However, there is one situation where we expect that the EMP may be useful, that is, for calculating properties of a system at a quantum critical point. At this point the correlations, which are due to entanglement, are strongest. We conjecture that the EMP (unlike the DMRG) may be able to reproduce the algebraic decay of correlations at the critical point, but have not yet been able to verify this analytically or numerically.

The difference between the entanglement preserved by the EMP and the DMRG projection may be seen using the example of the 1D antiferromagnetic Heisenberg model $H = J/2 \sum_{\langle i,j \rangle} \boldsymbol{\sigma}^i \cdot \boldsymbol{\sigma}^j$ (See [5, 21, 22, 23, 24] for other computations of the entanglement in condensed matter systems.) As we noted earlier, for computational ease we chose the model to be on four sites. The EMP was calculated by optimizing over the entanglement of formation $E(A : B)$ of the projected system using the formula of Wootters [25]. The entanglement of formation as a function of temperature is shown in Fig. 1. It is clear from Fig. 1 that the EMP better preserves the entanglement present in the system than does the DMRG, as expected. Similar results have been obtained for the Ising model in a transverse field, and we expect that this is typical of strongly interacting quantum lattice systems.

In this paper we have shown that entanglement plays a central role in quantum phase transitions. We have shown that the success of the DMRG — one of the principal numerical techniques used to solve quantum lattice systems near criticality — is due to the way it takes account of the entanglement in such systems. We have also




FIG. 1: Sites 1, 2 in the 4-site Heisenberg model were renormalized to a single spin via the DMRG and EMP projections (similarly for sites 3, 4) and the entanglement between the two renormalized spins plotted. Also shown is a lower bound on the actual entanglement present between A and B obtained from the entropy-entanglement inequality (chapter 8 in [5], and [20]). The lower bound is expected to be a good approximation only when $kT < 1$, where we see that the EMP provides a much better approximation to the entanglement present in the system than does the DMRG.

proposed a modification of the DMRG which preserves more entanglement during the renormalization step, and which we conjecture may be capable of reproducing the polynomial decay of correlations near the critical point. We believe our results show that the techniques of quantum information science can be fruitfully applied to obtain fundamental insights into the properties of many-body systems.

We thank Dorit Aharonov and John Preskill for stimulating and encouraging discussions about entanglement and phase transitions, and Jennifer Dodd for helpful conversations which substantially clarified the manuscript. This work was supported by an Australian Postgraduate Award to TJO.

* tjo@maths.uq.edu.au

† nielsen@physics.uq.edu.au

- [1] E. Schrödinger, Proc. Cambridge Phil. Soc. **31**, 555 (1935).
- [2] J. S. Bell, Physics **1**, 195 (1964).
- [3] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, Phys. Rev. A (3) **54**, 3824 (1996).
- [4] D. Aharonov, Phys. Rev. A **62**, 2311 (1999).
- [5] M. A. Nielsen, Ph.D. thesis, University of New Mexico (1998), quant-ph/0011036.
- [6] J. Preskill, J. Mod. Opt. **47**, 127 (2000).
- [7] S. Sachdev, *Quantum phase transitions* (Cambridge Uni-

- versity Press, Cambridge, 1999).
- [8] R. M. Noack and S. R. White, in *Density-matrix renormalization (Dresden, 1998)* (Springer, Berlin, 1999).
 - [9] K. G. Wilson, Rev. Modern Phys. **47**, 773 (1975).
 - [10] S. R. White, Phys. Rev. Lett. **69**, 2863 (1992).
 - [11] M. A. Nielsen and I. L. Chuang, *Quantum computation and quantum information* (Cambridge University Press, Cambridge, 2000).
 - [12] J. Preskill (1998), Physics 229 Lecture Notes online at <http://www.theory.caltech.edu/people/preskill/ph229/>.
 - [13] M. E. Peskin and D. V. Schroeder, *An introduction to quantum field theory* (Addison-Wesley Publishing Company Advanced Book Program, Reading, MA, 1995).
 - [14] M. E. Fisher, Rev. Modern Phys. **70** (1998).
 - [15] R. Bhatia, *Matrix analysis* (Springer-Verlag, New York, 1997).
 - [16] G. Vidal, D. Jonathan, and M. A. Nielsen, Phys. Rev. A (3) **62**, 012304 (2000).
 - [17] H. Barnum (1999), quant-ph/9910072.
 - [18] T. J. Osborne and M. A. Nielsen (2002), quant-ph/0202162.
 - [19] S. Rommer and S. Östlund, in *Density-matrix renormalization (Dresden, 1998)* (Springer, Berlin, 1999), pp. 67–89.
 - [20] P. Horodecki, R. Horodecki, and M. Horodecki, Acta Phys. Slov. **48**, 141 (1998).
 - [21] X. Wang, H. Fu, and A. I. Solomon (2001), quant-ph/0105075.
 - [22] D. Gunlycke, S. Bose, V. M. Kendon, and V. Vedral (2001), quant-ph/0102137.
 - [23] M. C. Arnesen, S. Bose, and V. Vedral, Phys. Rev. Lett. **87**, 2245 (2001).
 - [24] D. A. Meyer and N. R. Wallach (2001), quant-ph/0108104.
 - [25] W. K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998).